

ORIGINAL RESEARCH

# Effect of health literacy and socioeconomic deprivation on outcomes after lower limb surgical revascularisation for chronic limb-threatening ischaemia: the HeALTHI study

Bishop C,<sup>1</sup> El-Sayed T,<sup>1</sup> Baljer B,<sup>1</sup> Buckley E,<sup>1</sup> Convill J,<sup>1</sup> Rowlands G,<sup>2</sup> Bell R,<sup>1</sup> Nandhra S,<sup>1,2</sup>

1. The Northern Vascular Centre,  
Freeman Hospital,  
Newcastle upon Tyne, UK

2. Population Health Sciences  
Institute, Faculty of Medical  
Sciences,  
Newcastle University,  
Newcastle upon Tyne, UK

**Corresponding author:**

Tamer El-Sayed  
NIHR Clinical Lecturer in  
Vascular Surgery, The Northern  
Vascular Centre, Freeman  
Hospital, Freeman Road,  
High Heaton,  
Newcastle upon Tyne  
NE7 7DN, UK  
Email: [Tamer.El-Sayed@nhs.net](mailto:Tamer.El-Sayed@nhs.net)

**Received:** 23rd February 2023

**Accepted:** 5th October 2023

**Online:** 8th November 2023

## Plain English Summary

**Why we undertook the work:** Health literacy refers to the ability of patients to understand information related to their health condition so that they can make informed decisions about their health. Moreover, it is also believed that patients from disadvantaged backgrounds who have poor health are more likely to suffer from negative health outcomes. Therefore, we wanted to know whether this is true among our patients with poor leg circulation, particularly in its severe form which is called chronic limb-threatening ischaemia (CLTI) when they are undergoing surgery to improve their leg circulation.

**What we did:** We looked at patients who underwent major surgery for this disease (CLTI) in the Northern Vascular centre, Freeman Hospital, Newcastle upon Tyne, UK, and we asked them questions to see how much they understand about their condition, and we worked out how much advantaged or disadvantaged background they came from to know if that had any effect on how they did after surgery.

**What we found:** Our study found that a significant portion of patients with CLTI lacked knowledge about their condition and lived in poor areas. Unfortunately, we discovered that patients who lacked knowledge about their condition and hailed from poor areas were more likely to have complications after surgery.

**What this means:** Our research found that a patient's understanding of their health condition and the financial and social environment they live in can impact their recovery after surgery to improve their poor leg circulation. However, more studies are needed to better understand this relationship with a larger sample of patients.

## Abstract

**Introduction:** Research has suggested a relationship between health literacy, socioeconomic status and health-related outcomes. The aim of the study was to study the association between health literacy, socioeconomic status and outcomes following infra-inguinal bypass surgery for chronic limb-threatening ischaemia (CLTI).

**Methods:** Patients with CLTI undergoing lower limb surgical bypass graft operations between January 2016 and December 2018 were included in a cross-sectional observational study. The HLS19-Q12 questionnaire categorised participant's health literacy as inadequate, problematic, sufficient or excellent. Socioeconomic status was assessed using the Index of Multiple Deprivation (IMD). Primary outcomes were major lower limb amputation (MLLA) and adverse cardiovascular events. Secondary outcomes included length of hospital stay, and early postoperative complications including pneumonia, surgical site and graft infection. Kaplan–Meier survival curves were used to compare health literacy and amputation, and Cox proportional regression analysis was conducted to identify differences in limb loss risk against health literacy and social deprivation levels.

**Results:** The study consisted of 50 patients with an average age of 70±8.7 years. The participants' levels of health literacy were classified as inadequate (28%), problematic (38%), sufficient (24%) or excellent (10%). Approximately 40% of the patients lived in the most deprived areas. While all health literacy groups had similar postoperative outcomes, low health

literacy was connected with lower socioeconomic status ( $r=0.308$ ,  $p=0.029$ ). IMD ( $p=0.017$ , HR 0.502 (95% CI 0.285 to 0.883)) and haemoglobin ( $p=0.001$ , HR 0.919 (95% CI 0.872 to 0.968)) were significant predictors of MLLA.

**Conclusion:** Patients with lower health literacy are more likely to face higher levels of social deprivation, which may predict amputation following bypass surgery. Enhancing health literacy could play a role in reducing health disparities caused by social deprivation, thereby potentially addressing a vascular James Lind Alliance priority.

**Key words:** health literacy, social deprivation, health inequalities, chronic limb-threatening ischaemia

## Introduction

Health literacy is the degree to which individuals have the capacity to obtain, process and understand basic health information and access services needed to make appropriate health decisions.<sup>1</sup> Health literacy is a complex concept that is influenced by an individual's educational attainment, race, age, deprivation and available healthcare services.<sup>2,3</sup> There are a number of health literacy assessment tools available, including the Rapid Estimate of Adult Literacy in Medicine (REALM) test, the Test of Functional Health Literacy in Adults (TOFHLA), the Newest Vital Sign (NVS) and the 12-item European Health Literacy Survey (HLS19-Q12), which is a brief self-administered questionnaire that was developed by the European Health Literacy Survey (HLS) Consortium and been validated as a reliable measure of health literacy.<sup>4</sup>

Social deprivation has been associated with low health literacy, and is defined as the lack of access to the resources and opportunities that are considered to be essential for individuals to participate fully in society.<sup>1,5</sup> There are a number of social deprivation assessment tools available, including the Townsend deprivation index and the Carstairs deprivation score, which are based on census data, while the Index of Multiple Deprivation (IMD) is a composite measure of income, employment, education, housing and health.

Socioeconomic status is a social determinant of health and individuals with lower socioeconomic status have poorer health outcomes. Data support the association between social deprivation and health inequalities.<sup>6</sup> Social deprivation and health inequalities are particularly pertinent in vascular patients because vascular disease is strongly related to lifestyle through behaviours such as smoking, poor diet and lack of physical activity.<sup>7,8</sup> Chronic limb-threatening ischaemia (CLTI) is the severe presentation of peripheral arterial disease (PAD), producing ischaemic rest pain and tissue loss, reduced quality of life, and high risk of limb loss and/or mortality as well as increased cost and demand on vascular services.<sup>9</sup> Low health literacy can compromise a patient's adherence to lifestyle modification advice, medication compliance and the ability to seek medical advice when required.<sup>1,5</sup> Unfortunately, these are essentials in the management of PAD including CLTI. Inadequate health literacy was doubled among socially deprived cardiovascular patients compared with high socioeconomic status patients. These patients had lower rates of

access to health services, late presentation and poorer clinical outcomes.<sup>10-12</sup> Previous reports suggested that socially deprived vascular patients are at an increased risk of amputation up to 65% compared with the least deprived populations in the UK and the USA.<sup>13-15</sup>

Social deprivation and poor health behaviours not only contribute to disease development, but also to clinical outcomes after vascular intervention.<sup>7,16-18</sup> For example, peri-revascularisation smoking increases the risk of early major revascularisation bypass graft failure and limb loss.<sup>19,20</sup> Therefore, social deprivation is associated with an increased risk of major amputation following revascularisation, in contrast to patients with income above the poverty line.<sup>21-23</sup> A recent study suggested that over three-quarters of vascular patients had inadequate health literacy,<sup>24</sup> with an increased risk of hospital readmission, cardiovascular disease and mortality.<sup>25-28</sup> Therefore, understanding the relationship between health literacy and CLTI post-revascularisation outcomes is important given that health literacy is a potentially modifiable factor.<sup>1</sup> Therefore, there is a need for a better understanding of health literacy in this vascular patient group.

This study aimed to assess the relationship between health literacy, social deprivation and clinical outcomes in CLTI patients who have undergone revascularisation surgery.

The study primary research question was: Is health literacy associated with postoperative clinical outcomes in patients with CLTI following lower limb revascularisation surgery? Secondary research questions were: (1) Is there an association between health literacy score and social deprivation score in CLTI patients? and (2) Is the social deprivation score associated with poor clinical outcomes in patients with CLTI following lower limb revascularisation surgery?

## Methods

### Study design and population

This is a single-centre cross-sectional prospective study of all patients aged >18 years who underwent infra-inguinal lower limb bypass graft surgery for CLTI between January 2016 and December 2018 in the Northern Vascular Centre, Freeman Hospital, Newcastle upon Tyne. CLTI was defined by at least two weeks of ischaemic rest pain, lower limb ulceration or gangrene.

Health Research Authority and Research Ethics Committee

approvals were granted (21/NI/0092) with subsequent Newcastle upon Tyne Hospital research and development department approvals. Informed consent was obtained from all participants.

### Procedural data

All patients with CLTI included in the study underwent infra-inguinal lower limb bypass surgery with either autologous vein or prosthetic graft. Patients who underwent endovascular interventions only were not included in the study as primary endovascular procedures are mainly done as day case procedures which makes recruiting and assessing their health literacy more challenging compared with open/hybrid interventions, considering the intervention technicalities have little influence or relevance to the study question. All patients had their regular clinical follow-up as well as research team follow-up with median follow-up of 12 months.

### Baseline data collection

Baseline characteristics including demographics and comorbidities were collected by the research team from the patients' electronic healthcare records. Laboratory tests collected include preoperative haemoglobin, sodium, creatinine and estimated glomerular filtration rate.

Comorbidities are defined as per the Society of Vascular Surgery (SVS) guidelines where possible.<sup>29</sup> Diabetes is defined by documented medical history, the use of oral antidiabetic agents or insulin or fasting plasma glucose levels of at least 1.26 g/L; hypertension is defined by documented medical history and use of antihypertensive drugs for this purpose, or systolic blood pressure of at least 140 mmHg or diastolic blood pressure of at least 90 mmHg at admission determined by the average of the first two measurements. The following diseases were recorded based on the patient's documented medical history: ischaemic heart disease or prior myocardial infarction, atrial fibrillation, hypertension, cerebrovascular disease (stroke, including ischaemic or haemorrhagic stroke as well as transient ischaemic attack), end-stage renal failure, renal failure defined as estimated glomerular filtration rate of <15 mL/min/1.73 m<sup>2</sup> and/or requiring dialysis, and/or its documentation in their health records and a documented diagnosis of chronic obstructive pulmonary disease.


### Inclusion criteria (Figure 1)

- Patients aged  $\geq 18$  years.
- Patients who have undergone lower limb revascularisation surgery for CLTI at Newcastle upon Tyne Hospitals NHS Foundation Trust between 1 January 2015 and 1 January 2019.
- Patients able to consent to study participation.
- Patients able to answer the health literacy questionnaire.
- Patients who can speak/understand English.

### Exclusion criteria (Figure 1)

- Patients aged <18 years.
- Patients who have had lower limb revascularisation surgery for

**Figure 1** Consort of participant recruitment.



any reason other than CLTI.

- Patients who have had endovascular intervention even for CLTI.
- Patients who have had redo lower limb intervention even for CLTI.
- Patients who are unwilling to consent to study participation.
- Patients with incomplete data sets on the Northern Vascular Centre Register.
- Patients who are unable to speak/understand English.

### Socioeconomic status

Socioeconomic status was determined using the Office for National Statistics English Indices of Multiple Deprivation (IMD) 2019 measure.<sup>30</sup> This is a measure of relative deprivation and is calculated for each Lower-layer Super Output Area in England. The IMD tool graded participants' postcode areas, ranking them from 1 (most deprived) to 10 (least deprived).

### Health literacy

Health literacy was assessed using the 12-item European Health Literacy Survey (HLS19-Q12).<sup>4</sup> Each response on the HLS19-Q12 was assigned a numerical value ranging from 1 to 4. Subsequently, a total score ranging from 1 to 50 was calculated for each patient, following the HLS19 scoring system. Based on these scores, patients' health literacy was classified into four categories: inadequate (score 0–25 points), which is the severe level of poor health literacy; problematic (score >25–33 points), which is the moderate level of poor health literacy; sufficient (score >33–42 points), which is the mild level of poor health literacy; and excellent (score >42–50 points).

Postoperative outcomes encompassed major lower limb amputation (below, through or above the knee joint) and early postoperative complications including myocardial infarction, hospital acquired pneumonia, 3-month graft occlusion (first postoperative graft surveillance ultrasonography scan), surgical site infection (when documented in the patients' hospital/healthcare records and defined as an infection that occurs at the site of the

bypass and associated with cellulitis and/or purulent discharge) and raised inflammatory markers (leucocyte count and/or C reactive protein level).<sup>31</sup> Graft infection was diagnosed when documented in the patients' hospital/healthcare records and associated with overlying cellulitis, the presence of an exposed prosthetic graft, sinus tract with persistent purulent drainage and/or bleeding, or palpable anastomotic pseudoaneurysm. This is in addition to elevated inflammatory markers such as leucocyte count and/or C-reactive protein levels, and radiological evidence such as peri-graft fluid and/or gas, graft disruption and pseudoaneurysm formation observed on ultrasonography and computed tomography images.<sup>32</sup>

### Study outcomes

The primary outcome was to identify the association between health literacy and postoperative outcomes following lower limb revascularisation surgery for CLTI. Secondary outcomes included: (1) to identify if socioeconomic status was associated with an increased risk of limb loss (amputation-free survival) or mortality following lower limb revascularisation surgery for CLTI; and (2) to understand the relationship between socioeconomic status and health literacy scores.

### Statistical analyses

Normally distributed data were presented as mean (SD) and hypothesis testing was performed with paired and unpaired t-tests. Categorical data were analysed by means of a  $\chi^2$  test. A generalised linear model/Kruskal–Wallis test was used for comparison of mean/median continuous data between groups. IMD quintiles and health literacy categories were treated as ordinal and analysed with rank correlation. Spearman's rank correlation test was performed between IMD and health literacy. Kaplan–Meier survival curves were used with a log-rank test to compare health literacy and amputation. A Cox proportional regression analysis was conducted to identify differences in limb loss to enable hazard analysis. An a priori power analysis was conducted using G\*Power3 to test the primary outcomes and survival analyses, with an alpha of 0.05.<sup>33</sup> The results showed that a minimum sample of 48 and 35 participants were required, respectively, to achieve a power of 90%. A p value of <0.05 was considered statistically significant. Statistical analysis was performed using SPSS version 24 (IBM SPSS Statistics for Windows, Version 24.0. IBM Corp, Armonk, New York, USA).

### Results

A total of 50 participants were included in the study (39 males, 78.0%) with a mean±SD age of 70±8.7

years and median follow-up of 12 months. Fourteen patients (28%) had inadequate health literacy, 19 (38%) had problematic health literacy, 12 (24%) had sufficient health literacy and 5 (10%) had excellent health literacy. Twenty-five participants (50%) lived in areas of highest deprivation (IMD 1–3), 11 participants (22%) lived in areas of moderate deprivation (IMD 4–6) and 14 (28%) lived in areas of lowest deprivation (IMD 7–10). Baseline demographics were comparable between health literacy groups (Table 1).

A significant difference was found between IMD groups based on the degree of severity of lower limb ischaemia as outlined by the Rutherford grade at the time of intervention (p=0.029) and hypertension (p=0.031) (Table 2). Health literacy was weakly but significantly correlated with IMD score (r=0.308, p=0.029).

There were no significant differences in postoperative outcomes between health literacy groups (Table 3) or IMD quintiles (Table 4), including length of stay, 3-month graft patency, MLLA and myocardial infarction.

The mortality rate over the 12-month period according to health literacy and IMD groups was as follows: Health literacy: inadequate 21.42%, problematic 15.78%, sufficient 8.33%, excellent 0.00%.

**Table 1** Baseline characteristics by health literacy group.

Variable	Health literacy group				P value
	Inadequate (n=14)	Problematic (n=19)	Sufficient (n=12)	Excellent (n=5)	
Male	12	14	10	3	
Age, years	71 (7)	71 (9)	70 (10)	66 (10)	
IMD quintile					0.124
1	6	8	5	1	
2	4	3	3	0	
3	2	3	1	0	
4	2	3	1	2	
5	0	2	2	2	
Comorbidities					
Diabetes mellitus	9	6	5	1	0.201
Ischaemic heart disease	4	6	6	1	0.657
Hypertension	8	12	6	3	0.619
Cardiac failure	2	0	0	0	0.157
Renal failure	0	1	0	0	0.613
Chronic respiratory disease	1	4	2	1	0.698
Pre-surgery medication					
Statins	12	12	10	4	0.723
Single antiplatelet	10	12	10	4	0.834
Rutherford grade					0.512
4	4	9	5	2	
5	2	4	0	1	
6	7	4	5	2	
Missing data (n=5)	1	2	2	–	
Laboratory tests					
Haemoglobin, g/L	127 (23)	133 (23)	126 (18)	129 (9)	0.813
Leukocytes, $\times 10^3$ mean	9.40 (1.91)	8.63 (2.70)	9.76 (3.08)	8.52 (2.70)	0.646
Total protein, g/L	70 (7)	64 (9)	67 (8)	76 (11)	0.399
Creatinine, $\mu\text{mol/L}$	89(23)	83 (29)	105 (97)	76 (17)	0.712

Continuous data are presented as mean (SD).  
IMD, Indices of Multiple Deprivation.

**Table 2** Baseline characteristics by IMD quintile

Variable	IMD Quintile					P value
	1 (n=20)	2 (n=10)	3 (n=6)	4 (n=8)	5 (n=6)	
Male	15	7	5	7	5	
Age, years	69 (9)	69 (6)	68 (8)	71(13)	77 (8)	0.330
Health literacy						
Inadequate	6	4	2	2	0	0.124
Problematic	8	3	3	3	2	
Sufficient	5	3	1	1	2	
Excellent	1	0	0	2	2	
Comorbidities						
Diabetic	8	4	3	5	1	0.525
Ischaemic heart disease	5	2	4	5	1	0.064
Hypertension	13	3	2	8	3	0.031
Cardiac failure	0	1	0	1	0	0.492
Renal failure	0	0	0	1	0	0.321
Chronic respiratory disease	4	3	0	0	1	0.349
Pre-surgery medication						
Statins	14	8	5	7	4	0.565
Single antiplatelet	13	8	5	6	4	0.682
Rutherford grade						0.029
4	6	6	1	6	1	0.221
5	2	2	1	1	1	
6	10	1	3	1	3	
Missing data	2	1	1	–	1	
Laboratory tests						
Haemoglobin, g/L						0.808
Leukocytes, x10 <sup>3</sup>	9.56 (2.17)	8.11 (1.42)	10.63 (3.09)	8.77 (2.57)	8.70 (4.72)	0.408
Total protein, g/L						
Creatinine, µmol/L	95 (80)	80 (27)	87 (14)	94 (35)	90 (25)	0.978

Continuous data are presented as mean (SD). IMD, Indices of Multiple Deprivation.

**Table 3** Lower limb revascularisation outcomes by health literacy group

Outcomes	Health literacy group				P value
	Inadequate (n=14)	Problematic (n=19)	Sufficient (n=12)	Excellent (n=5)	
Length of stay (days)	25 (22)	12(15)	19 (25)	7(5)	0.201
3-month graft patency	3	8	1	1	0.157
MLLA	5	5	4	0	0.487
MI	0	1	0	0	0.683
Wound infection	1	3	2	0	0.715
Graft infection	1	0	2	0	0.277

Continuous data are presented as mean (SD). MLLA, major lower limb amputation; MI, myocardial infarction.

IMD: 1st quintile 30%, 2nd quintile 10%, 3rd quintile 33.3%, 4th quintile 12.5%, 5th quintile 0.00%. Kaplan–Meier analysis revealed no significant difference in amputation-free survival between health literacy groups (log rank  $p=0.545$ ) and IMD groups (log rank  $p=0.887$ ) (Figure 2A). Cox regression analysis showed that IMD ( $p=0.017$ , HR 0.502 (95% CI 0.285 to 0.883)) and haemoglobin ( $p=0.001$ , HR 0.919 (95% CI 0.872 to 0.968)) were significant

predictors of MLLA (Table 5). Only one patient developed postoperative myocardial infarction, so Cox regression could not be calculated for this outcome.

## Discussion

To our knowledge, this is the first study to investigate the association between health literacy among patients with CLTI undergoing lower limb revascularisation bypass surgery and postoperative outcomes. We found that socioeconomic status is associated with the health literacy level and the degree of severity of vascular disease in patients who underwent vascular intervention for CLTI. Social deprivation level measured by IMD was found to be a significant predictor of major adverse limb events; however, there was no significant difference in major adverse clinical outcomes among health literacy groups.

Our study cohort showed a weak correlation between health literacy and IMD. This correlation was also demonstrated by a previous report and showed that lower socioeconomic groups are more likely to have low health literacy, resulting in poor health status. This is because health literacy was shown to mediate the relationship between socioeconomic status and health status, quality of life and health-related outcomes. This could be particularly true in patients with PAD as their socioeconomic status is likely to influence their health behaviours such as smoking, poor diet and physical activity.<sup>34,35</sup>

Approximately half of our study population were from areas of high levels of social deprivation. Patients from high levels of social deprivation were likely to have inadequate health literacy, as a significant correlation was found between socioeconomic status and health literacy ( $p=0.029$ ). This was better explained when 70% of participants in the lowest IMD quintile had inadequate or problematic health literacy, while in the highest IMD quintile only 33% had problematic health

literacy. This is comparable to the European study which found limited health literacy in 73.9% of individuals with low social status.<sup>36</sup>

Although no difference was found in postoperative outcomes between IMD groups, similar to a UK multiple national databases study, there was no association between patients' socioeconomic deprivation and one-year postoperative death following major amputation for CLTI, but the authors concluded that the relationship



**Table 4** Lower limb revascularisation outcomes by IMD quintile

Variable	IMD Quintile					P value
	1 (n=20)	2 (n=10)	3 (n=8)	4 (n=6)	5 (n=8)	
Length of stay, days	22 (25)	12 (14)	23 (19)	10 (13)	10 (10)	0.460
3-month graft patency	5	0	4	3	1	0.057
MLLA	7	2	3	2	0	0.348
MI	1	0	0	0	0	0.852
Wound infection	2	1	1	0	2	0.520
Graft infection	2	1	0	0	0	0.736


Continuous data are presented as mean (SD).

IMD, Indices of Multiple Deprivation; MLLA, major lower limb amputation; MI, myocardial infarction.


between social deprivation and CLTI is potentially more complex and suggested a prospective investigation of this relationship.<sup>37</sup> However, our Cox regression analysis showed that IMD was a significant predictor of major lower limb amputation. This finding is supported by previous studies that showed patients from deprived areas were 2.4 times more at risk of amputation compared with less deprived areas.<sup>38</sup>

In our cohort, although not statistically significant, participants from areas of highest deprivation were younger (69 years) compared with the participants in the lowest deprivation


**Figure 2** Amputation-free survival predictors. (A) Kaplan–Meier amputation-free survival curve by health literacy group (log rank  $p=0.54$ ). Vertical lines indicate censored data. (B) Kaplan–Meier amputation-free survival curve by IMD quintiles ( $p=0.017$ ). (C and D) Cox regression hazard plots showing MLLA risk as per IMD ( $p=0.044$ ) and anaemia ( $p=0.001$ ).




Patients at risk	0	1	2	3	4	5	6
Inadequate	14	11	10	10	6	4	2
Problematic	19	15	15	14	9	4	2
Sufficient	12	11	10	10	9	8	1
Excellent	5	5	4	4	3	1	1



Patients at risk	0	1	2	3	4	5	6
1st Quintile	20	14	14	13	10	6	2
2nd Quintile	10	9	9	9	4	3	1
3rd Quintile	6	4	3	3	3	2	1
4th Quintile	8	7	6	5	2	1	1
5th Quintile	6	6	6	6	6	4	1



Patients at risk	0	1	2	3	4	5	6
1st Quintile	20	14	14	13	10	6	2
2nd Quintile	10	9	9	9	4	3	1
3rd Quintile	6	4	3	3	3	2	1
4th Quintile	8	7	6	5	2	1	1
5th Quintile	6	6	6	6	6	4	1



Patients at risk	0	1	2	3	4	5	6
Non-anaemic	33	29	28	26	18	13	4
Anaemic	13	9	8	8	5	3	1

IMD, Indices of Multiple Deprivation (Quintile); MLLA, major lower limb amputation.

**Table 5** Predictors of major lower limb amputation following revascularisation by Cox regression analysis

	Hazard ratio	95% CI	P value
IMD	0.502	0.285 to 0.883	0.017
HL	0.799	0.344 to 1.860	0.603
Age	1.047	0.955 to 1.147	0.311
Sex	0.368	0.050 to 2.690	0.324
Diabetes mellitus	2.665	0.558 to 12.740	0.219
Ischaemic heart disease	1.104	0.361 to 3.377	0.863
Hypertension	0.688	0.128 to 3.701	0.663
COPD	0.252	0.026 to 2.405	0.231
Rutherford grade	1.330	0.444 to 3.984	0.611
Haemoglobin	0.919	0.872 to 0.968	0.001

COPD, chronic obstructive pulmonary disease; HL, health literacy; IMD, Indices of Multiple Deprivation (Quintile).

areas (77 years). Previous reports have found individuals from high deprivation areas presented with similar pathology and comorbidities at a younger age compared with their less socially deprived counterparts.<sup>39</sup>

In our patient cohort, none of the participants with excellent health literacy had an amputation. One could assume that excellent health literacy could be a protective effect against major amputation. Nevertheless, caution is advised as the 'excellent health literacy group' was small in this study ( $n=5$ ), and a larger cohort of patients in a wider study is needed for better evaluation of this association.

The majority of our participants (66%) had inadequate or problematic health literacy, which is considerably higher than the 47.6% of individuals with inadequate or problematic health literacy in a previous European study.<sup>31</sup> Similar high rates of low health literacy (76.7%) were reported among PAD clinic patients in the Netherlands.<sup>21</sup> Limited data exist for a comparison of health literacy in England. Although our cohort of patients did not show a significant difference in postoperative outcomes among health literacy groups, other surgical areas have found significant differences. For example, patients with poor health literacy spent a longer time in the hospital following major abdominal surgery and suffered from an increased risk of complications such as surgical site infections and, when they were discharged, they had decreased compliance with discharge instructions including medication compliance and wound/drain care.<sup>40–42</sup>

Social deprivation combined with reduced health literacy can be linked with an increased display of poor lifestyle behaviours including smoking, poor diet and inadequate physical activity leading to poorer health outcomes. This could be particularly true in patients with PAD as their socioeconomic status is likely to influence their health behaviours such as smoking, poor diet and physical activity, which are common risk factors for PAD and amputation.<sup>8,43–45</sup>

Lack of awareness of vascular disease symptoms and inequalities in access to healthcare services may lead to socially deprived patients presenting with more advanced vascular disease at a younger age. Requiring surgical intervention at a younger age could result in higher disability life-years experienced and more life-years lost, therefore leading to reduced health-related quality of life.<sup>39,46</sup>

Recent studies have shown that patients with adequate health literacy will have the knowledge, skills and confidence to navigate the healthcare system, leading to more efficient use of healthcare services and better health outcomes.<sup>35,47</sup> Improving health literacy could therefore have a mediating role in the relationship between social deprivation and health outcomes through health literacy interventions. This could improve the outcomes for patients with PAD following vascular intervention.

In this study we have shown that health literacy could be a social determinant of health, as health literacy follows a social gradient and creates health inequalities. We believe health literacy is a potentially modifiable factor and could be a facilitator in reducing the gap in health inequalities and social deprivation, with subsequent improved health outcomes.<sup>35,47</sup> Health literacy intervention remains in its infancy, with most research being conducted in the USA, including patient-centred communication and self-management programmes.<sup>48,49</sup>

Health literacy is influenced by race and ethnicity as individuals from different cultures may struggle to navigate through the healthcare system due, for example, to language barrier, different levels of education and different beliefs about health and illness. A number of studies have shown that people from racial and ethnic minority groups are more likely to have low health literacy than those from white majority groups. For example, a study by the National Centre for Education Statistics found that 44% of African Americans and 42% of Hispanics had low health literacy compared with 28% of white subjects.<sup>50</sup>

Improving health literacy is recognised as a national imperative, prompting collaborative efforts between Health Education England, Public Health England and NHS England, and it was one of the foremost priorities identified by the James Lind Alliance for PAD management coupled with improving the educational support for patients with poor leg circulation. Therefore, health literacy enhancement could be a potential avenue for mitigating health inequities stemming from socioeconomic disadvantages, thereby aligning with a key vascular priority established by the James Lind Alliance.<sup>51</sup> Health literacy interventions should be tailored at an individual level for an improved patients' understanding and more efficient use of healthcare services, better health outcomes and improved quality of life for patients.<sup>2,3</sup> As social deprivation and health literacy are linked, there may be merit in identifying vascular patients with a low IMD postcode and targeting these patients for perioperative risk modification to improve postoperative outcomes.<sup>39</sup>

## KEY MESSAGES

- There is an association between poor health literacy and social deprivation among patients with CLTI.
- Patients with poor health literacy coming from socially deprived areas are at higher risk of amputation following lower limb revascularisation for CLTI.

## Study limitations

The study has several limitations that should be acknowledged. First, the sample size was relatively small, consisting of patients from a single centre. Additionally, the study population was relatively homogenous, comprising individuals residing in the North-East of England. Therefore, the findings may not be generalisable to broader populations. Despite these limitations, the study provides valuable insights and warrants future investigation through larger multicentre prospective observational studies. These studies should aim to evaluate the impact of health literacy and social deprivation on patients with CLTI and their revascularisation outcomes.

## Conclusion

This study provides valuable insights into the clinical implications of health literacy and social deprivation in patients with PAD. There is an association between poor health literacy and social deprivation among patients with CLTI. Patients with poor health literacy coming from socially deprived areas are at higher risk of MLLA following lower limb revascularisation for CLTI. Based on the findings of this study, tailored health literacy interventions could be implemented to mitigate the negative impact of low health literacy and social deprivation on postoperative outcomes.

**Conflict of Interest:** None.

**Funding:** None.

**Acknowledgement:** The HLS19 instrument used in this research was developed within "HLS19 – the International Health Literacy Population Survey 2019–2021" of M-POHL. The authors would like to thank the M-POHL Network for kindly giving us permission to use their HLS19-Q12 questionnaire.

**Reviewer acknowledgement:** *JVSGBI* thanks Bharad Ravindhran, Hull University Hospitals NHS Trust and Andrew Nickinson, University Hospitals of Southampton NHS Foundation Trust, for their contribution to the peer review of this work.

## References

- Rowlands G, Tabassum B, Campbell P, *et al*. The evidence-based development of an intervention to improve clinical health literacy practice. *Int J Environ Res Public Health* 2020;**17**(5):1513. <https://doi.org/10.3390/ijerph17051513>
- Dunn P, Conard S. Improving health literacy in patients with chronic conditions: a call to action. *Int J Cardiol* 2018;**273**:249–51. <https://doi.org/10.1016/j.ijcard.2018.08.090>
- Rowlands G, Protheroe J, Winkley J, Richardson M, Seed PT, Rudd R. A mismatch between population health literacy and the complexity of health information: an observational study. *Br J Gen Pract* 2015;**65**(635):e379–e386. <https://doi.org/10.3399/bjgp15X685285>
- The HLS19 Consortium of the WHO Action Network M-POHL. International report on the methodology, results, and recommendations of the European health literacy. Population survey 2019–2021 (HLS19) of M-POHL. Austrian National Public Health Institute, Vienna, 2021. <https://m-pohl.net/sites/m-pohl.net/files/inline-files/HLS19%20International%20Report.pdf>
- Suhail M, Saeed H, Saleem Z, *et al*. Association of health literacy and medication adherence with health-related quality of life (HRQoL) in patients with ischemic heart disease. *Health Qual Life Outcomes* 2021;**19**(1):118. <https://doi.org/10.1186/s12955-021-01761-5>
- Marmot M. Health equity in England: the Marmot review 10 years on. *BMJ* 2020;**368**:m693. <https://doi.org/10.1136/bmj.m693>
- Conte MS, Bradbury AW, Kolh P, *et al*. Global vascular guidelines on the management of chronic limb-threatening ischemia. *Eur J Vasc Endovasc Surg* 2019;**58**(1S):S1–S109.e33. [Published correction appears in *Eur J Vasc Endovasc Surg* 2020;**59**(3):492–3; published correction appears in *Eur J Vasc Endovasc Surg* 2020;**60**(1):158–9]. <https://doi.org/10.1016/j.ejvs.2019.05.006>
- Pujades-Rodriguez M, Timmis A, Stogiannis D, *et al*. Socioeconomic deprivation and the incidence of 12 cardiovascular diseases in 1.9 million women and men: implications for risk prediction and prevention. *PLoS One* 2014;**9**(8):e104671. <https://doi.org/10.1371/journal.pone.0104671>
- Ying AF, Tang TY, Jin A, Chong TT, Hausenloy DJ, Koh WP. Diabetes and other vascular risk factors in association with the risk of lower extremity amputation in chronic limb-threatening ischemia: a prospective cohort study. *Cardiovasc Diabetol* 2022;**21**(1):7. <https://doi.org/10.1186/s12933-021-01441-0>
- Diederichs C, Jordan S, Domanska O, Neuhauser H. Health literacy in men and women with cardiovascular diseases and its association with the use of health care services - results from the population-based GEDA2014/2015-EHIS survey in Germany. *PLoS One* 2018;**13**(12):e0208303. <https://doi.org/10.1371/journal.pone.0208303>
- Wittink H, Oosterhaven J. Patient education and health literacy. *Musculoskeletal Sci Pract* 2018;**38**:120–7. <https://doi.org/10.1016/j.msksp.2018.06.004>
- Eronen J, Paakkari L, Portegijs E, Saajanaho M, Rantanen T. Health literacy supports active aging. *Prev Med* 2021;**143**:106330. <https://doi.org/10.1016/j.ypmed.2020.106330>
- Venermo M, Manderbacka K, Ikonen T, Keskimäki I, Winell K, Sund R. Amputations and socioeconomic position among persons with diabetes mellitus, a population-based register study. *BMJ Open* 2013;**3**(4):e002395. <https://doi.org/10.1136/bmjopen-2012-002395>
- Ferguson HJ, Nightingale P, Pathak R, Jayatunga AP. The influence of socio-economic deprivation on rates of major lower limb amputation secondary to peripheral arterial disease. *Eur J Vasc Endovasc Surg* 2010;**40**(1):76–80. <https://doi.org/10.1016/j.ejvs.2010.03.008>
- Stevens CD, Schriger DL, Raffetto B, Davis AC, Zingmond D, Roby DH. Geographic clustering of diabetic lower-extremity amputations in low-income regions of California. *Health Aff (Millwood)* 2014;**33**(8):1383–90. <https://doi.org/10.1377/hlthaff.2014.0148>
- Li X, Galvin JW, Li C, Agrawal R, Curry EJ. The impact of socioeconomic status on outcomes in orthopaedic surgery. *J Bone Joint Surg Am* 2020;**102**(5):428–44. <https://doi.org/10.2106/JBJS.19.00504>
- Pagano D, Freemantle N, Bridgewater B, *et al*. Social deprivation and prognostic benefits of cardiac surgery: observational study of 44 902 patients from five hospitals over 10 years. *BMJ* 2009;338:b902. <https://doi.org/10.1136/bmj.b902>
- Mirza AH, Aylin P, Middleton S, King EV, Nouraei RAR, Repanos C. Impact of social deprivation on the outcome of major head and neck cancer surgery in England: a national analysis. *Head Neck* 2019;**41**(3):692–700. <https://doi.org/10.1002/hed.25461>
- Kalbaugh CA, Gonzalez NJ, Luckett DJ, *et al*. The impact of current smoking on outcomes after infrainguinal bypass for claudication. *J Vasc Surg* 2018;**68**(2):495–502.e1. <https://doi.org/10.1016/j.jvs.2017.10.091>
- Selvarajah S, Black JH 3rd, Malas MB, Lum YW, Propper BW, Abularrage CJ. Preoperative smoking is associated with early graft failure after infrainguinal bypass surgery. *J Vasc Surg* 2014;**59**(5):1308–14. <https://doi.org/10.1016/j.jvs.2013.12.011>
- Heikkilä K, Loftus IM, Waton S, Johal AS, Boyle JR, Cromwell DA. Association of neighbourhood deprivation with risks of major amputation and death following lower limb revascularisation. *Atherosclerosis* 2020;**306**:11–14. <https://doi.org/10.1016/j.atherosclerosis.2020.06.017>
- Gurney JK, Stanley J, York S, Rosenbaum D, Sarfati D. Risk of lower limb amputation in a national prevalent cohort of patients with diabetes. *Diabetologia* 2018;**61**(3):626–35. <https://doi.org/10.1007/s00125-017-4488-8>
- Durham CA, Mohr MC, Parker FM, Bogey WM, Powell CS, Stoner MC. The impact of socioeconomic factors on outcome and hospital costs associated with femoropopliteal revascularization. *J Vasc Surg* 2010;**52**(3):600–7. <https://doi.org/10.1016/j.jvs.2010.04.011>
- Srijbos RM, Hinnen JW, van den Haak RFF, Verhoeven BAN, Koning OHJ.



- Inadequate health literacy in patients with arterial vascular disease. *Eur J Vasc Endovasc Surg* 2018;**56**(2):239–45. <https://doi.org/10.1016/j.ejvs.2018.04.015>
25. Lindahl B, Norberg M, Johansson H, *et al*. Health literacy is independently and inversely associated with carotid artery plaques and cardiovascular risk. *Eur J Prev Cardiol* 2020;**27**(2):209–15. <https://doi.org/10.1177/2047487319882821>
  26. Uemura K, Yamada M, Kuzuya M, Okamoto H. The association of limited health literacy and risk of arterial stiffness in community-dwelling older adults [in Japanese]. *Nihon Ronen Igakkai Zasshi* 2018;**55**(4):605–11. <https://doi.org/10.3143/geriatrics.55.605>
  27. Baker S, Malone E, Graham L, *et al*. Patient-reported health literacy scores are associated with readmissions following surgery. *Am J Surg* 2020;**220**(5):1138–44. <https://doi.org/10.1016/j.amjsurg.2020.06.071>
  28. Mayberry LS, Schildcrout JS, Wallston KA, *et al*. Health literacy and 1-year mortality: mechanisms of association in adults hospitalized for cardiovascular disease. *Mayo Clin Proc* 2018;**93**(12):1728–38. <https://doi.org/10.1016/j.mayocp.2018.07.024>
  29. Writing Committee to Develop Clinical Data Standards for Peripheral Atherosclerotic Vascular Disease; Creager MA, Belkin M, Bluth EI, *et al*. 2012 ACCF/AHA/ACR/SCAI/SIR/STS/SVM/SVN/SVS key data elements and definitions for peripheral atherosclerotic vascular disease: a report of the American College of Cardiology Foundation/American Heart Association Task Force on Clinical Data Standards. *Circulation* 2012;**125**(2):395–467. [Erratum published in *Circulation* 2012;**125**(7):e387–8. Erratum published in *Circulation* 2013;**128**(11):e179]. <https://doi.org/10.1161/CIR.0b013e31823299a1>
  30. Ministry of Housing, Communities and Local Government. English indices of deprivation 2019. HM Government, 2019. Available from: <https://imd-by-post-code.opendatacommunities.org/imd/2019>.
  31. Horan TC, Andrus M, Dudeck MA. CDC/NHSN surveillance definition of health care-associated infection and criteria for specific types of infections in the acute care setting. *Am J Infect Control* 2008;**36**:309–32. <https://doi.org/10.1016/j.ajic.2008.03.002>
  32. Vicaretti M. Pathophysiology of vascular graft infections. In: Fitridge R, Thompson M, editors. *Mechanisms of Vascular Disease: A Reference Book for Vascular Specialists* [Internet]. Adelaide: University of Adelaide Press, 2011, p. 29. <https://www.ncbi.nlm.nih.gov/books/NBK534262/>
  33. Faul F, Erdfelder E, Lang AG, Buchner A. G\*Power 3: a flexible statistical power analysis program for the social, behavioral, and biomedical sciences. *Behav Res Methods* 2007;**39**(2):175–91. <https://doi.org/10.3758/bf03193146>
  34. Gibney S, Bruton L, Ryan C, Doyle G, Rowlands G. Increasing health literacy may reduce health inequalities: evidence from a national population survey in Ireland. *Int J Environ Res Public Health* 2020;**17**(16):5891. <https://doi.org/10.3390/ijerph17165891>
  35. Stormacq C, Van den Broucke S, Wosinski J. Does health literacy mediate the relationship between socioeconomic status and health disparities? Integrative review. *Health Promot Int* 2019;**34**(5):e1–e17. <https://doi.org/10.1093/heapro/day062>
  36. Sørensen K, Pelikan JM, Röthlin F, *et al*. Health literacy in Europe: comparative results of the European health literacy survey (HLS-EU). *Eur J Public Health* 2015;**25**(6):1053–8. <https://doi.org/10.1093/eurpub/ckv043>
  37. Nickinson ATO, Coles B, Payne TJ, Davies RSM, Khunti K, Sayers RD. Is socio-economic deprivation associated with one year post-operative mortality following major amputation for chronic limb threatening ischaemia? *Eur J Vasc Endovasc Surg* 2021;**61**(6):1026–7. <https://doi.org/10.1016/j.ejvs.2021.02.020>
  38. Hicks KA, Tchong JE, Bozkurt B, *et al*. 2014 ACC/AHA key data elements and definitions for cardiovascular endpoint events in clinical trials: a report of the American College of Cardiology/American Heart Association Task Force on Clinical Data Standards (Writing Committee to Develop Cardiovascular Endpoints Data Standards). *J Am Coll Cardiol* 2015;**66**(4):4036–9. [Published correction appears in *J Am Coll Cardiol* 2015;**66**(8):982]. <https://doi.org/10.1016/j.jacc.2014.12.018>
  39. Gwilym BL, Maheswaran R, Edwards A, *et al*. Income deprivation and groin wound surgical site infection: cross-sectional analysis from the groin wound infection after vascular exposure multicenter cohort study. *Surg Infect (Larchmt)* 2022;**23**(1):73–83. <https://doi.org/10.1089/sur.2021.153>
  40. Theiss LM, Wood T, McLeod MC, *et al*. The association of health literacy and postoperative complications after colorectal surgery: a cohort study. *Am J Surg* 2022;**223**(6):1047–52. <https://doi.org/10.1016/j.amjsurg.2021.10.024>
  41. Calabrese EC, Asmar S, Bible L, *et al*. Prospective evaluation of health literacy and its impact on outcomes in emergency general surgery. *J Surg Res* 2021;**261**:343–50. <https://doi.org/10.1016/j.jss.2020.12.047>
  42. Wright JP, Edwards GC, Goggins K, *et al*. Association of health literacy with postoperative outcomes in patients undergoing major abdominal surgery. *JAMA Surg* 2018;**153**(2):137–42. <https://doi.org/10.1001/jamasurg.2017.3832>
  43. Jayasinghe UW, Harris MF, Parker SM, *et al*. The impact of health literacy and life style risk factors on health-related quality of life of Australian patients. *Health Qual Life Outcomes* 2016;**14**:68. <https://doi.org/10.1186/s12955-016-0471-1>
  44. Feinglass J, Shively VP, Martin GJ, *et al*. How 'preventable' are lower extremity amputations? A qualitative study of patient perceptions of precipitating factors. *Disabil Rehabil* 2012;**34**(25):2158–65. <https://doi.org/10.3109/09638288.2012.677936>
  45. RobatSarpoooshi D, Mahdizadeh M, Alizadeh Siuki H, Haddadi M, Robatsarpoooshi H, Peyman N. The relationship between health literacy level and self-care behaviors in patients with diabetes. *Patient Relat Outcome Meas* 2020;**11**:129–35. <https://doi.org/10.2147/PROM.S243678>
  46. Rocha V, Ribeiro AI, Severo M, Barros H, Fraga S. Neighbourhood socioeconomic deprivation and health-related quality of life: a multilevel analysis. *PLoS One* 2017;**12**(12):e0188736. <https://doi.org/10.1371/journal.pone.0188736>
  47. Pelikan JM, Ganahl K, Roethlin F. Health literacy as a determinant, mediator and/or moderator of health: empirical models using the European Health Literacy Survey dataset. *Glob Health Promot* 2018;**25**(4):1757975918788300. <https://doi.org/10.1177/1757975918788300>
  48. Sudore RL, Schillinger D. Interventions to improve care for patients with limited health literacy. *J Clin Outcomes Manag* 2009;**16**(1):20–9.
  49. Visscher BB, Steunenberg B, Heijmans M, *et al*. Evidence on the effectiveness of health literacy interventions in the EU: a systematic review. *BMC Public Health* 2018;**18**(1):1414. <https://doi.org/10.1186/s12889-018-6331-7>
  50. Muvuka B, Combs RM, Ayangeakaa SD, Ali NM, Wendel ML, Jackson T. Health literacy in African-American communities: barriers and strategies. *Health Lit Res Pract* 2020;**4**(3):e138–e143. <https://doi.org/10.3928/24748307-20200617-01>
  51. Pymer S, Harwood AE, Long J, Chetter IC, Saratzis A, Coughlin P, on behalf of the Vascular Society of Great Britain and Ireland Peripheral Arterial Disease Special Interest Group James Lind Alliance Priority Setting Partnership. Research priorities for patients with peripheral arterial disease: a James Lind Alliance Priority Setting Partnership. *J Vasc Soc GB Irel* 2022;**1**(2):23–9. <https://doi.org/10.54522/jvsobi.2022.011>