

CASE REPORT

Rare vascular complication of total hip arthroplasty: common femoral artery perforation diagnosed two months postoperatively

Mouhanni S, Mouyarden O, Farah S, El Jamaaoui A, El Bhali H, Azghari A

Department of Vascular Surgery,
University Hospital of Tangier,
University Abdelmalek Essaadi,
Tangier, Morocco

Corresponding author:
Dr Ouassim Mouyarden
Department of Vascular Surgery,
University Hospital of Tangier,
University Abdelmalek Essaadi,
BP 917, Tangier, Morocco
Email: o.mouyarden@gmail.com

Received: 15th August 2025
Accepted: 24th October 2025
Online: 17th November 2025

Abstract

Background: Vascular injuries complicating total hip arthroplasty (THA) are uncommon, with reported incidences between 0.2% and 0.42%. When they occur, they can threaten both the patient's life and functional prognosis of the limb.

Case report: We describe a 38-year-old woman with tuberculous coxitis who underwent THA. Two months later she presented with acute back and groin pain associated with severe anaemia. Exploration revealed a perforation of the common femoral artery (CFA). Reconstruction using reversed saphenous vein bypass and profunda femoris reimplantation resulted in successful recovery.

Conclusion: CFA perforation after THA is exceedingly rare and may present late. Any unexplained groin symptoms or anaemia following THA should trigger urgent vascular evaluation.

Key words: hip arthroplasty; vascular injury; common femoral artery; pseudo-aneurysm; vascular surgery

Introduction

Arterial complications following total hip arthroplasty (THA) are rare but potentially devastating, with incidence estimated between 0.2% and 0.42%.^{1,2} Mechanisms include penetrating trauma from retractors or screws, traction injury during dislocation, perforation during acetabular preparation, or delayed vessel erosion leading to pseudoaneurysm formation.³ Revision surgery, dysplasia and pelvic fractures are established risk factors.⁴

We report an exceptional case of delayed common femoral artery (CFA) perforation two

months after primary uncomplicated THA for tuberculous coxitis, highlighting the need for vigilance beyond the immediate postoperative period.

Case report

Preoperative assessment

A 38-year-old woman with treated pulmonary tuberculosis presented with an 18-month history of right hip pain and limping. Examination revealed restricted flexion without vascular or neurological deficits. A pelvic radiograph demonstrated destruction of the femoral head and acetabular changes consistent with chronic coxitis (Figure 1). A CT scan confirmed chronic right-sided coxitis with muscular atrophy (Figure 2). Bone biopsy verified tuberculous infection. After multidisciplinary review she underwent THA using the Hardinge lateral approach. Intraoperative fluoroscopy was satisfactory and the early postoperative course was uneventful (Figure 3), with postoperative haemoglobin 9.6 g/dL.

Postoperative presentation

Two months later the patient developed sudden low back and groin pain radiating to the thigh. The hip was held in flexion; distal pulses remained palpable. Laboratory results showed severe anaemia (haemoglobin 4 g/dL). Because she was haemodynamically stable, surgical exploration was performed without preoperative angiography.

Operative findings and management

A large haematoma was evacuated, revealing active arterial bleeding at the femoral bifurcation. The vascular team was called. After proximal and distal control, a transfixing perforation of the CFA was identified (Figures 4 and 5). The damaged

Figure 1 Preoperative pelvic radiograph showing a deformed right femoral head with osteolysis and joint space narrowing.




Figure 2 CT scan images of the right hip demonstrating chronic coxitis and acetabular destruction.




Figure 3 Postoperative pelvic radiograph after total hip arthroplasty.




Figure 4 Intraoperative photograph showing the common femoral artery perforation at the femoral bifurcation.




Figure 5 Intraoperative photograph showing control of the femoral bifurcation.





Figure 6 Intraoperative photograph of the venous bypass graft with reimplantation of the profunda femoris artery.



segment was excised and continuity restored with a reversed autologous saphenous vein graft from the CFA to the superficial femoral artery, with profunda femoris re-implanted into the graft (Figure 6). The patient recovered uneventfully with preserved limb perfusion.

KEY MESSAGES

- Vascular injury following THA is rare but serious. Published series report a 0.2–0.42% incidence of arterial injury. Low incidence contributes to delayed recognition.
- Delayed presentation is possible. CFA perforation or pseudoaneurysm may present weeks or months after surgery with groin pain, anaemia or neurological symptoms even when distal pulses are preserved.
- Maintain a high index of suspicion. Any unexplained groin, thigh or back pain, swelling or drop in haemoglobin after THA should trigger vascular evaluation.
- Early multidisciplinary management saves limbs. Prompt recognition and collaboration between orthopaedic and vascular surgeons allow timely repair and minimise morbidity.
- Meticulous surgical technique and preoperative planning reduce risk. Awareness of anatomical variations, careful placement of retractors and screws, and avoidance of excessive traction or torsion are essential preventive measures.

Discussion

Vascular injury during THA remains rare yet potentially catastrophic. Reported incidence ranges between 0.2% and 0.42%.¹ The external iliac and common femoral arteries are most often involved because of their proximity to the acetabulum and femoral canal.² Early recognition is crucial, since delayed diagnosis can lead to limb loss or death.³

Mechanisms include direct trauma from retractors, drills or acetabular screws; thermal damage from bone cement; traction-related intimal disruption; and delayed perforation from pseudoaneurysm erosion.^{4,5} Our patient lacked traditional risk factors – no revision surgery, dysplasia, pelvic fracture or fibrosis – emphasising that even technically uncomplicated primary THA can produce vascular complications.

Delayed presentations, such as pseudoaneurysm rupture months after surgery, are diagnostically challenging. Collateral circulation may maintain distal pulses, masking acute ischaemia.⁶ Therefore, persistent groin pain, thigh swelling, neurological deficit or unexplained anaemia after THA should raise suspicion for vascular injury. CT angiography remains the diagnostic modality of choice when haemodynamic stability allows.

Management depends on lesion type and location. Endovascular therapy is increasingly preferred for contained perforations or pseudoaneurysms in anatomically suitable sites.⁷ However, open reconstruction remains essential when the femoral bifurcation is involved, when infection is possible, or when prosthetic artefacts limit imaging accuracy.⁸ Autologous saphenous vein provides durable infection-resistant reconstruction, as illustrated in this case.

This experience underlines the necessity of continued postoperative vigilance and close collaboration between orthopaedic and vascular teams. Even in primary THA, awareness of vascular anatomy, meticulous technique and prompt multidisciplinary action are vital to preventing devastating outcomes.

Conclusion

Common femoral artery perforation following THA is exceptionally rare but potentially life-threatening. Delayed presentation, as in this case, complicates diagnosis and management. Persistent groin pain, anaemia or neurological symptoms after hip arthroplasty should always prompt vascular evaluation. Early diagnosis and multidisciplinary collaboration between orthopaedic and vascular teams remain vital for limb salvage and survival. Preventive strategies include meticulous surgical technique, respect for anatomic variations and careful postoperative vigilance.

Conflict of Interest: None.

Funding: None.

Patient consent to publication: Written informed consent was obtained from the patient for publication of this case report and any accompanying images.

References

1. Calligaro KD, Dougherty MJ, Ryan S, Booth RE. Acute arterial complications associated with total hip and knee arthroplasty. *J Vasc Surg* 2003;**38**:1170–7. [https://doi.org/10.1016/s0741-5214\(03\)00918-2](https://doi.org/10.1016/s0741-5214(03)00918-2)
2. Abularrage CJ, Weiswasser JM, Dezee KJ, Slidell MB, Henderson WG, Sidawy AN. Predictors of lower-extremity arterial injury after total knee or total hip arthroplasty. *J Vasc Surg* 2008;**47**(4):803–7. <https://doi.org/10.1016/j.jvs.2007.11.067>
3. Street MWJ, Howard LC, Neufeld ME, Masri BA. Vascular injuries during hip and knee replacement. *Orthop Clin North Am* 2022;**53**(3):395–408. <https://doi.org/10.1016/j.ocl.2021.08.009>
4. Meknarit S, Motta JC, de Grandis E, W Anthony Lee. Diagnosis and management of vascular injuries after joint arthroplasty. *Annals of Vascular Surgery* 2023;**3**(2):100199. <https://doi.org/10.1016/j.av surg.2023.100199>
5. Mohan V, Gopal VS. Delayed vascular complication after total hip replacement. *Journal of Orthopaedic Association of South Indian States* 2022;**19**(2):95–8. https://doi.org/10.4103/joasis.joasis_23_22
6. Katira K, Martin A, Garbuzov A, *et al.* Peripheral arterial disease and complications of total knee arthroplasty: indications for advanced vascular imaging and minimally invasive soft-tissue coverage procedures. *Orthoplastic Surgery* 2023;**14**:1–8. <https://doi.org/10.1016/j.orthop.2023.09.002>
7. Proschek D, Proschek P, Hochmuth K *et al.* False aneurysm of the left femoral artery and thrombosis of the left femoral vein after total hip arthroplasty. *Arch Orthop Trauma Surg* 2006;**126**:493–7. <https://doi.org/10.1007/s00402-006-0166-x>
8. Al-Salman M, Taylor DC, Beauchamp CP, Duncan CP. Prevention of vascular injuries in revision total hip replacement. *Can J Surg* 1992;**35**:261–4.